

Role of Clove in Human Medical History

Muhammad Arshad Ullah ^{1*} | Ali Hassan ^{2*} | Ameer Hamza ³

¹ Pakistan Agricultural Research Council, Islamabad, Pakistan

² PMAS- University of Arid Agriculture, Rawalpindi, Pakistan

³ COMSATS- Biosciences Department, Islamabad Campus, Pakistan

***Corresponding Author:** Ali Hassan, PMAS- University of Arid Agriculture, Rawalpindi, Pakistan

Received Date: July 15,2025; **Accepted Date:** July 28,2025; **Published Date:** August 02,2025

Citation: Muhammad Arshad Ullah, Ali Hassan, Ameer Hamza (2025). Role of Clove in Human Medical History, *J Role of Clove in Human Medical History*. 2(4) 28, **DOI:** BRCA-RW-26-28

Copyright: Ali Hassan, et al © (2025). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract:

The clove is usually known as "lavang." Due to the increase in nutrition and medicinal properties, Plants' importance in human life has increased every day. Clove is the topical evergreen Myrtaceae family tree native to the islands of Maluka in east Indonesia. It is commonly used for tooth care. Clove is used extensively in dental care for relieving toothache, sore gums and oral ulcers. Gargling with clove oil can also aid in sore throat conditions and bad breathe. This is commonly used as an expectorant for the treatment of dental discomfort, the digestive problem, oral ulceration, mosquito repellent. Clove is a pharmacologically active medicinal plant that includes Anti-oxidant, anti-pyretic, anti-viral, anti-microbial, anti-diabetic, anti-inflammatory, analgesic, anti-platelet, anti-stress, anti-disease, and anti-carcinogenic in cervical cancer. Clove is amongst the most essential sources of Phenolic compounds, including eugenol (80% -90%), eugenyl acetate (15% -17%), and beta-caryophyllene (5% -12%), alpha-humulene (0.55%), alpha-terpenyl acetate (0.1%), and methyl eugenol (0.2%). The clove oil extracted from the dried floral buds of the clove provide as a topical treating pain and as a healing factor. The flowers, stalks and clove tree leaves are obtained from the distillation of clove oil. Eugenol is the clove's principal bioactive ingredient. For oral diseases, it is used as an antiseptic.

Cloves are used in the cuisine of Asian, African, Middle East countries and used to give aromatic and flavor qualities to hot beverages. A mine component of clove plant parts is eugenol, that it has not been classified for its potential toxicity. Other important essential oil of clove is acetyl eugenol, beta-caryophyllene and vanillin, categolic acid, tannins such as bicornin, gallotannic acid, methyl salicylate, the flavonoids eugenin, kaempferol, rhamnetin, and eugenitin, triterpenoids such as oleanolic acid, stigmasterol, and campesterol and several sesquiterpenes. Thus, foods should be preserved against the microbial spoilage throughout the storage periods. In addition to clove extracts could affects as anti-oxidative, fungicidal, and antibacterial effects on foods. It has been reported that clove essential oil is one of the natural fungicides and antibacterial phytomaterial. The active essential oil in clove, eugenol, has been shown to act as an effective platelet inhibitor, preventing blood clots. The major constituents in bud and leaf oils were reported to be eugenol and caryophyllene. Main constituent's flower buds of clove essential oil are phenylpropanoids such as carvacrol, thymol, eugenol and cinnamaldehyde. Clove bud oil contained primarily eugenol, eugenyl acetate and caryophyllene. The high concentration of eugenol in leaf and buds oil makes it potentially useful in the medicines because they exhibit antibacterial, antifungal, anti-inflammatory activity, insecticidal and antioxidant properties, and are used traditionally as flavouring agent and antimicrobial material in food. Clove (*Syzygium aromaticum* L.) buds and flowers are utilized for its essential oil. Eugenol is the main component of clove oil, which has strong antioxidant activity. Eugenol is the chief bioactive constituent of clove, which is present in concentrations ranging from 9 381.70 to 14 650.00 mg/100 g of fresh plant weight. With regard to the phenolic acids, gallic acid is found in higher concentration (783.50 mg/100 g fresh weight).

Cloves contain appreciable amounts of volatile oil (used for flavouring foods and pharmaceuticals), which is mainly confined in aerial parts of plant. Other phenolic acids found in clove are caffeic, ferulic, elagic and salicylic acids. Flavonoids including kaempferol, quercetin and its derivates (glycosilated) are also found in trace amounts.

Appreciable amounts of essential oil are present in aerial parts of clove. Good quality clove bud contains volatile oil (15 to 20%), which mainly comprises of eugenol (70 to 85%), eugenyl acetate (10 to 15%), and beta-caryophyllene (5 to 12%). Approximately, 15-20% volatile oil can be produced from dried buds. Clove improves memory by relieving mental fog, drowsiness, and depression. Clove oil is mosquito repellent. Clove oil exhibited significant inhibition against tested strains, with minimum inhibitory concentrations ranging from 0.312-1.25% (v/v) for clove oil, and 0.312-5% (v/v) for rosemary oil.

Key words: clove; mosquito; memory

Introduction:

Clove is used extensively in dental care for relieving toothache, sore gums and oral ulcers. Gargling with clove oil can also aid in sore throat conditions and bad breathe. Arabic traders brought the buds to Europe in 4th century A.D., and in the seventh and eighth century A.D. Europe, cloves became very popular as a medicinal flower, due to their ability to preserve foods, and mask the smell of poorly-kept foods (International Organization for Standardization, 2002).

The active essential oil in clove, eugenol, has been shown to act as an effective platelet inhibitor, preventing blood clots (Körfers *et al.*, 2009).

Preliminary studies suggest that clove oil may play a chemo preventive role, particularly in cases of lung, skin and digestive cancers (Schönfelder *et al.*, 2004).

Clove bud oil has biological activities, such as antibacterial, antifungal, insecticidal and antioxidant properties, and are used traditionally as flavouring agent and antimicrobial material in food (Lee and Shibamoto, 2001; Huang *et al.*, 2002; Velluti *et al.*, 2003).

Importantly, Lee and Shibamoto, 2001, reported that clove oil might also be used as an anti-carcinogenic agent due to its antioxidant properties. Their results also suggested that clove oil might be of use as a potential chemo preventative agent.

Syzygium species have been reported to possess antibacterial (Shafi *et al.*, 2002) and anti-inflammatory activity (Muruganadan *et al.*, 2001).

Recently, flavonoid triglycosides have been isolated (Nassar, 2006). The major constituents in bud and leaf oils were reported to be eugenol and caryophyllene (Srivastava *et al.*, 2003; Raina *et al.*, 2001; Wenqiang *et al.*, 2007). Kamel *et al.* (2007) reported the main constituent's flower buds of clove essential oil are phenylpropanoids such as carvacrol, thymol, eugenol and cinnamaldehyde. Amla *et al.* (2007) also reported that the bud oil contained primarily eugenol, eugenyl acetate and -caryophyllene.

The high concentration of eugenol in leaf and buds oil makes it potentially useful in the medicines because they exhibit antibacterial, antifungal, anti-inflammatory activity, insecticidal and antioxidant properties, and are used traditionally as flavouring agent and antimicrobial material in food (Huang *et al.*, 2002; Velluti *et al.*, 2003; Shafi *et al.*, 2002; Muruganadan *et al.*, 2001).

Clove (*Syzygium aromaticum* L.) buds and flowers are utilized for its essential oil. In Indonesia, clove is classified into four main varieties: Siputih, Sikotok, Zanzibar, and Ambon (Luthfi, 2017). The Zanzibar variety produces more buds than other varieties and considered the most superior and cultivated variety (Rukmana and Yudirachman, 2016). A clove bud is a single flower with a length of 1-2 cm, while a clove flower has eight stages of clove bud development, namely young bud, budding-1, budding-2, budding-3, full-budding, flowering, initial fruiting, and full fruiting (Rukmana and Yudirachman, 2016 and Razafimamponjison *et al.*, 2013). Eugenol is the main component of clove oil, which has strong antioxidant activity (Laitupa and Susane, 2010).

There are obvious physiological differences between unstable and stable growth stage. However, studies on its yield and quality as essential oil are limited. Particularly, young trees at 3-4 years old had the first flowering period (Semangun and Karwur, 2014), which makes the flower buds an interesting material for experiment.

The highest oil yield in young trees was also reported in several previous studies on other plant species (Pujiarti *et al.*, 2011 and Pambudi *et al.*, 2016). Composition of the vacuole structures led to high oil yield in young clove trees. Essential oil is one of the secondary metabolites stored in plant vacuoles (Campbell and Reece, 2012). Small vacuoles merge to form a single large structure as the plants grow (Evert, 2006).

The biosynthesis starts from the first budding stage, meaning the transformation of the droplet into bubble occurs along with the budding stages (Kridati *et al.*, 2012).

Optical rotation of oil extracted from young trees was higher than that from mature trees, which was similar to that found in eucalyptus oil (Runtunuwu *et al.*, 2016). According to Pujiarti *et al.* (2011), tree age has a diverse influence on the optical rotation of essential oils.

Changes in the polarization angle increase with the decrease in the viscosity of the material (Nuraniza *et al.*, 2013).

The clove is usually known as "lavang." Due to the increase in nutrition and medicinal properties, Plants' importance in human life has increased every day. The term clove is derived from 'clove' and 'clou' meaning 'nail.' This is used mainly in Ayurvedics (Hussain and Trak, 2019). Clove is the topical evergreen Myrtaceae family tree native to the islands of Maluka in east Indonesia.

The color is small brown, floral buds, and a spice is used. It was used as food preservatives, flavoring agents and nutritional additives, medicinal coloring agents (Cortés-Rojas *et al.*, 2014). Health, cosmetic, nutritional and agricultural use clove. It is commonly used for tooth care (Chaieb *et al.*, 2007). This is commonly used as an expectorant for the treatment of dental discomfort, the digestive problem, oral ulceration, mosquito repellent (Thangaselvabai *et al.*, 2010). Clove is a pharmacologically active medicinal plant that includes Anti-oxidant, anti-pyretic, anti-viral, anti-microbial, anti-diabetic, anti-inflammatory, analgesic, anti-platelet, anti-stress, anti-disease, and anti-carcinogenic in cervical cancer. The clove is available in three types of essential oil (Jain, 2019). Clove is amongst the most essential sources of Phenolic compounds, including eugenol (80% -90%), eugenyl acetate (15% -17%), and beta-caryophyllene (5% -12%), alpha-humulene (0.55%), alpha-terpenyl acetate (0.1%), and methyl eugenol (0.2%) (Yunusa *et al.*, 2018). The clove oil extracted from the dried floral buds of the clove provide as a topical treating pain and as a healing factor. The flowers, stalks and clove tree leaves are obtained from the distillation of clove oil (Khalil *et al.*, 2017). Eugenol is the clove's principal bioactive ingredient. For oral diseases, it is used as an antiseptic (Overly, 2019). The clove tree is an evergreen plant. Cloves flower consist of a long calyx that terminates in four spreading sepals, and four unopened petals that form a small central ball with aromatic scent (Bhuiyan *et al.*, 2010). The original clove zone is in Indonesia and Philippines (Danthu *et al.*, 2014). Cloves are used in the cuisine of Asian, African, Middle East countries and may be used to give aromatic and flavor qualities to hot beverages. A mine component of clove plant parts is eugenol, that it has not been classified for its potential toxicity (Bendre *et al.*, 2016). Other important essential oil of clove is acetyl eugenol, beta-caryophyllene and vanillin, crategolic acid, tannins such as bicornin, gallotannic acid, methyl salicylate, the flavonoids eugenin, kaempferol, rhamnetin, and eugenitin, triterpenoids such as oleanolic acid, stigmasterol, and campesterol and several sesquiterpenes (Bendre *et al.*, 2016). Thus, foods should be preserved against the microbial spoilage throughout the storage periods (Goni *et al.*, 2009).

In addition to clove extracts could affects as anti-oxidative, fungicidal, and antibacterial effects on foods (Goni *et al.*, 2009). It has been reported that clove essential oil is one of the natural fungicides and antibacterial phytomaterial (Ayoola *et al.*, 2008). The garden strawberry need to remain on the plant to fully ripen because they do not continue to ripen after being picked (Raina *et al.*, 2001). Rotted and overripe berries are removed to minimize insect and disease problems (Sallato *et al.*, 2007). The berries do not get washed until just before consumption. In large operations, strawberries are fall victim to a number of diseases like powdery mildew, leaf spot, leaf blight and slime molds (Pleasant, 2011).

The use of biocontrol agents with plant extracts in plant disease control has been employed by Zeng *et al.* (2012) suggested that clove extract might be a viable alternative to synthetic fungicides to extend the postharvest storage period and maintain fruit quality of navel orange too.

Weight loss occurred during storage regardless the type of fruit or vegetable evaluated. However, the rate of water loss was dependent on the type of crop evaluated, and was greatly related to the physiological and morphological characteristics of each individual fruit or vegetable, and with of each individual fruit or vegetable, and with the expected shelf life under the environmental conditions used in this study (Pino *et al.*, 2001 and Tournas and Katsoudas, 2005).

Clove enhances memory retention. It is recommended for relieving brain fog, lethargy and depressive state of mind. Research has shown that clove oil is an effective mosquito repellent (Trongtokit *et al.*, 2005).

Clove oil showed antimicrobial activity against some human pathogenic bacteria resistant to certain antibiotics (Lopez *et al.*, 2005). Clove oil and its main component eugenol show considerable antifungal activity against *Candida Aspergillus* and dermatophyte species. It also shows activity against clinically relevant fungi including fluconazole- resistant strains (Pinto *et al.*, 2009).

Clove is a potent antiviral agent. Eugenin isolated from clove buds showed antiviral activity against Herpes Simplex virus at a concentration of 10 μ g /ml (Chaieb *et al.*, 2007). It also down regulates the expression of some growth promoting proteins, viz, COX², cMyc, Hras (Banerjee *et al.*, 2006).

Oral administration of aqueous infusions of clove at the dose of 100 μ l /mouse/day not only delayed the formation of papilloma but also reduced the incidence of papilloma as well as the cumulative number of papillomas per mouse (Banerjee *et al.*, 2005). Clove and Eugenol possess strong antioxidant activity, which is comparable to the activities of the synthetic antioxidant, BHA (butylated hydroxyl anisole) and Pyrogallol (Dorman *et al.*, 2000). It also showed a significant inhibitory effect against hydroxyl radicals and act as an iron chelator (Gulcin *et al.*, 2004).

Cloves showed the highest DPPH radical scavenging activity & highest FRAP values (Yadav and Bhatnagar, 2007). The antioxidant activity of clove bud extract and its major aroma components, eugenol and eugenol acetate were comparable to that of the natural antioxidant α -tocopherol (Lee and Shibamoto, 2001). Eugenol inhibited 5-lipoxygenase activity and leukotriene C-4 in human PMNL cells (Raghavendra *et al.*, 2006).

A more global analysis of gene expression by DNA microarray analysis revealed that clove and insulin regulated the expression of many of the same genes in a similar manner (Prasad *et al.*, 2005).

Clove also contains a variety of flavonoids, including kaempferol, rhamnetin and β -caryophyllene which also contributed to clove's anti -inflammatory and antioxidant properties (Ghelardini *et al.*, 2001).

Clove extract was also effective in increasing the latency of anoxic stress induced convulsions in mice (Singh *et al.*, 2009). It has been found that ethanolic extract of clove (50%) produced a significant and sustained increase in the sexual activity of normal male rats, without any conspicuous gastric ulceration and adverse effects (Tajuddin *et al.*, 2004).

Culex quinquefasciatus and *Anopheles dirus* under laboratory conditions using human subjects (Trongtokit *et al.*, 2005).

The clover leaf and bud oils showed potent insecticidal activity against the human head louse (*Pediculus capitis*) (Yang *et al.*,

2003).

Clove in particular has attracted the attention due to the potent antioxidant and antimicrobial activities standing out among the other spices (Shan *et al.*, 2005). For centuries the trade of clove and the search of this valuable spice stimulated the economic development of this Asiatic region (Kamatou *et al.*, 2012). The collection could be done manually or chemically-mediated using a natural phytohormone which liberates ethylene in the vegetal tissue, producing precocious maturation (Filho *et al.*, 2013). Nowadays, the larger producer countries of clove are Indonesia, India, Malaysia, Sri Lanka, Madagascar and Tanzania specially the Zanzibar Island (Kamatou *et al.*, 2012). In Brazil, clove is cultured in the northeast region, in the state of Bahia in the regions of Valença, Ituberá, Taperoá, Camamu and Nilo Peçanha, where approximately 8 000 hectares are cultivated, producing near 2500 tons per year (Oliveira *et al.*, 2007 and 2009).

Concentrations up to 18% of essential oil can be found in the clove flower buds. Roughly, 89% of the clove essential oil is eugenol and 5% to 15% is eugenol acetate and β -cariofileno (Jirovetz *et al.*, 2006).

The antioxidant activity of aqueous extracts of clove has been tested by different in vitro methods as 2, 2-diphenyl-1-picrylhydrazyl (DPPH); 2, 2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) (ABTS), oxygen radical absorbance capacity, ferric reducing antioxidant power, xanthine oxidase and 2-deoxiguanosine. Clove and plants as pine, cinnamon, and mate proved its enormous potential as food preservative among the other 30 plants analyzed (Dudonné *et al.*, 2009). The powerful antioxidant activity of both extracts may be attributed to the strong hydrogen donating ability, metal chelating ability and scavenging of free radicals, hydrogen peroxide and superoxide (Gülçina *et al.*, 2004). Gülcin *et al.* (2011) studied the antioxidant activity of eugenol by several in vitro methods and discusses the structure-activity relationship (Gülçina *et al.*, 2011). Listeria monocitogenes, results reinforces the employment of eugenol to inhibit the growth of these microorganisms in surfaces in contact with food (Pérez-Conesa *et al.*, 2014).

Rana *et al.* determined the antifungal activity of clove oil in different strains and reported this scale of sensibility Mucor sp. > Microsporum gypseum > Fusarium moniliforme NCIM 1100 > Trichophyton rubrum > Aspergillus spp. > Fusarium oxysporum MTCC 284 (Rana *et al.*, 2011). The chromatographic analyses showed that eugenol was the main compound responsible for the antifungal activity due to lysis of the spores and micelles. A similar mechanism of action of membrane disruption and deformation of macromolecules produced by eugenol was reported by Devi *et al.* (2010).

The activities of clove oil against different dermatophytes as Microsporum canis (KCTC 6591), Trichophyton mentagrophytes (KCTC 6077), Trichophyton rubrum (KCCM60443), Epidermophyton floccosum (KCCM 11667) and Microsporum gypseum were tested and results indicate a maximum activity at concentration of 0.2 mg/mL with an effectiveness of up to 60% (Park *et al.*, 2007).

Pure clove oil or mixes with rosemary (*Rosmarinus officinalis* spp.) oil were tested against *Staphylococcus epidermidis*, *Staphylococcus aureus*, *Bacillus subtilis*, *E. coli*, *Proteus vulgaris*, *Pseudomonas aeruginosa* and results showed minimum inhibitory concentrations between 0.062% and 0.500% (v/v) which is promising as antiinfectious agents or as food preservative (Fu *et al.*, 2007).

The anticandidal activity of eugenol and carvacrol was tested in a vaginal candidiasis model, microbial and histological techniques were employed to compare the samples with the controls. The results suggest that eugenol and carvacrol could be a promising antifungal agent for treatment and prophylaxis of vaginal candidiasis (Chami *et al.*, 2004).

In addition to the wide spectrum of activity of eugenol against bacteria, a study showed that eugenol and cinnamaldehyde at 2 μ g/mL inhibited the growth of 31 strains of *Helicobacter pylori*, after 9h and 12h of incubation, respectively, being more potent than amoxicillin and without developing resistance. The activity and stability of those compounds was checked at low pH values since *Helicobacter pylori* resides in the stomach (Ali *et al.*, 2005).

The results showed the increase in the therapeutic effectiveness of eugenol and the modification of the release when administrated as solid lipid nanoparticles (Garg and Singh, 2011).

Beta-cyclodextrin inclusion complexes containing eugenol and clove bud extracts were tested against two common foodborne pathogens, *Salmonella enterica* var *Typhimurium* LT2 and *Listeria innocua* (Hill *et al.*, 2013).

The mechanism evolved has been attributed to the activation of calcium and chloride channels in ganglionar cells (Healthcare, 2004). The voltage dependant effects of eugenol in sodium and calcium channels and in receptors expressed in the trigeminal ganglio also contributed to the analgesic effect of clove (Li *et al.*, 2008). The peripheral antinociceptive activity of eugenol was reported by Daniel *et al.* showing significant activity at doses of 50, 75 and 100 mg/kg (Daniel *et al.*, 2009). This factor is activated by free radicals and results in the expression of genes that suppress apoptosis and induce cellular transformation, proliferation, invasion, metastasis among others (Aggarwal and Shishodia, 2006).

With regard to toxicity, eugenol presented strong genotoxic effects (DNA-damaging) on human VH10 fibroblast, medium genotoxic effects on Caco-2 colon cells and non DNA-damaging effects on HepG2 hepatome cells (Slamenová *et al.*, 2009). Nevertheless the National Toxicology Program based on several long term carcinogenicity studies concluded that eugenol was not carcinogenic to rats (Ghosh *et al.*, 2005). The cytotoxicity, reactive oxygen species (ROS) production, and intracellular glutathione levels in a human submandibular cell line (HSG cells) of eugenol and isoeugenol was studied by Atsumi *et al.* (2005). Therefore, it can be concluded that the cytotoxicity of eugenol occurs in a ROS-independent manner in the presence of oxidative stress. In another work, it was reported that eugenol inhibits the enzyme MMP-9 which is related to metastasis in human fibrosarcoma cells suggesting its application for prevention of metastasis related to oxidative stress (Nam and Kim, 2013).

Danio rerio and *Poecilia reticulata* the medium lethal concentrations (LD50) at 96h were (18.2-5.52) mg/mL in *Danio rerio* and (21.7-0.8) mg/mL in *Poecilia reticulata* (Doleželová *et al.*, 2011).

Eugenol is easily absorbed when administrated by oral route reaching rapidly plasma and blood with mean half-lives of 14.0h and 18.3h, respectively. Acumulative effect has been hypothesized and associated to relieve of neuropathic pain after repeated daily administrations (Guénette *et al.*, 2007).

Park and Shin (2005) reported the possibility of employment of clove essential oil to control the jponesse terminte Reticulitermes speratus (Kolbe *et al.*, 2009). In the same way, Eamsobhana *et al.* (2009) found that clove essencial oil at 5%posses 100%of repellent activity against the chigger Leptotrombidium imphalu which could be a safer and cheaper alternative to synthetic repellents commonly associated to harmful side effects (Sritabutra *et al.*, 2011).

The larvical methods are one of the most effective strategies to combat dengue, since there is not drug for treatment or a vaccine. Eugenol exhibited interesting results and could be a promising alternative to common insecticide (Barbosa *et al.*, 2013).Eugenol, eugenol acetate and beta-caryophyllene were effective in repellency of red imported fire ants *Solenopsis invicta* (Hymenoptera: Formicidae), being eugenol the fastest acting compound (Zhang *et al.*, 2013). Clove oil was also effective spatial repellent for pestiferous social wasps *Vespa pensylvanica* (Saussure) and paper wasps mainly *Polistes dominulus* (Christ) (Jawahery *et al.*, 2012).

Clove oil can also serve as an anesthesia for a variety of fish. However, lengthy exposures can cause mortality and sub-acute morbidity (Hekimoğlu and Ergun, 2012).

Clove oil could be employed as suppressor of potato tuber germination by affecting the lipid peroxidation and the enzymes activities of catalase, glutathione-S-transferase, peroxidase, polyphenol oxidase and superoxide dismutase (Afify *et al.*, 2012). Syzygium is the largest genus of Mirtaceae family, comprising of about 1200 to 1800 species of flowering plants, which are widely distributed in tropical and subtropical areas of Asia, Africa, Madagascar, and throughout Pacific and Oceanic regions (Cock and Cheesman, 2018).

Cloves contain appreciable amounts of volatile oil (used for flavouring foods and pharmaceuticals), which is mainly confined in aerial parts of plant. The yield and composition of volatile oil are variable and are thought to be linked to growing conditions, genetic factors, different chemotypes, geographic origins, and differences in the nutritional status of plant (Al-Maskri *et al.*, 2011; Arshad *et al.*, 2014 and Hanif *et al.*, 2019). Clove is known by different vernacular names in different languages. It is known as qaranful (Arabic), Karamfil (Bulgarian), Ding xiang (Chinese), Kruidnagel (Danish), Garifalo (Greek), Mikhaki (Georgian), Nelke (German), Szegfu (Hungarian), Cengkeh (Indonesian), Choji (Japanese), Jeonghyang (Korean), Krustnaglinas (Latvian), Lwaang (Nepalese), Carvo de India (Portuguese), Mikhak (Persian), Kala (Pashto), Gvosdika (Russian), Clavo (Spanish), Carenfil (Turkish), Garn ploo (Thai), Dhing huong (Vietnamese), and Laung (Urdu/Punjabi/Hindi) (Milind and Deepa, 2011).

Eugenol is the chief bioactive constituent of clove, which is present in concentrations ranging from 9 381.70 to 14 650.00 mg/100 g of fresh plant weight. With regard to the phenolic acids, gallic acid is found in higher concentration (783.50 mg/100 g fresh weight) (Shan *et al.*, 2005). Other phenolic acids found in clove are caffeic, ferulic, elagic and salicylic acids. Flavonoids including kaempferol, quercetin and its derivates (glycosilated) are also found in trace amounts. Appreciable amounts of essential oil are present in aerial parts of clove. Chemical profile of this oil is generally found by GCMS analysis (Ahmad *et al.*, 2008; Hanif *et al.*, 2011 and Javed *et al.*, 2012). Good quality clove bud contains volatile oil (15 to 20%), which mainly comprises of eugenol (70 to 85%), eugenyl acetate (10 to 15%), and beta-caryophyllene (5 to 12%). Other minor constituents including methyl amyl ketone, kaempferol, gallotannic acid, α -humulene, β -humulene, methyl salicylate, categolic acid, and benzaldehyde are responsible for the characteristic pleasant fragrance of clove (Mittal *et al.*, 2014).

Drying may take 4 to 5 days. Well dried buds are hard, crisp and dark brown, having moisture content (<12%), which can be stored for 1 to 2 years in gunny bags. Approximately, 15-20% volatile oil can be produced from dried buds (Thangaselvabai *et al.*, 2010).

Clove improves memory by relieving mental fog, drowsiness, and depression. Clove oil is mosquito repellent (Trongtokit *et al.*, 2005).

Clover oil exhibited significant inhibition against tested strains, with minimum inhibitory concentrations ranging from 0.312-1.25% (v/v) for clove oil, and 0.312- 5% (v/v) for rosemary oil (Abdullah *et al.*, 2015).

Clove oil was tested against five dermatophytes including *Trichophyton rubrum*, *Epidermophyton floccosum*, *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton mentagrophytes*. Maximum inhibitory effect (\approx 60%) against all fungal strains was shown at dose of 0.2mg/ml (Park *et al.*, 2007).

Results revealed the maximum inhibitory action of clove, mustard, and cinnamon at 1% concentration. Garlic showed good inhibitory action at 3% concentration. However, mint and ginger had negligible inhibition at same concentration (Sofia *et al.*, 2007). Essential oils of *Piper nigrum*, *Syzygium aromaticum*, *Pelargonium graveolens*, *Myristica fragrans*, *Origanum vulgare*, and *Thymus vulgaris* were evaluated for antimicrobial activity against twenty five bacterial strains, including food borne, animal, and plant pathogens considerable inhibitory action was observed by the volatile oils in a dose dependent behavior (Dorman and Deans, 2000).

It was also verified that the citotoxic potential of eugenol was more powerful than borneol. With regard to toxicity, eugenol exhibited strong DNA damaging effects on human fibroblast (VH10), medium damaging effects on colon cells (caco-2) and non genotoxic effects on hepatome cells (HepG2) (Slameňová *et al.*, 2009).

Generally phenolic and flavonoids are responsible for antioxidant activities of the oil (Khan *et al.*, 2012). Among all, clove showed maximum, whereas, onion showed minimum inhibitory potential (Shobana and Naidu, 2000).

Essential oils were added to soybean oil at doses of 0.006 and 0.01g/ml, for thirty days, at accelerated oxidation level. Among all examined oils, the clove oil showed more potent ($p<0.05$) antioxidant activity followed by oregano and sage oils (Ghadermazi

et al., 2017).

Eugenin isolated from clove bud essential oil exhibited a potent inhibitory effect against herpes simplex virus at a dose of 10 μ g/ml (Chaiet al., 2007).

The degree of hepatic damage was evaluated by increased levels of cytoplasmic enzymes (aspartate aminotransferase and alanine aminotransferase). Clove extract restored the normal concentrations of enzymes in serum (Thuwaini et al., 2016).

The toxicity of clove oil was evaluated in aquarium fish species, Poecilia reticulata and Danio rerio. The LD50 values were (18.2 \pm 5.52) mg/ml and (21.7 \pm 0.8) mg/ml in Danio rerio and Poecilia reticulata, respectively, at 96 h (Doleželová et al., 2011)

References:

1. Abdullah B.H., S.F. Hatem, W. Jumaa. (2015). A comparative study of the antibacterial activity of clove and rosemary essential oils on multidrug resistant bacteria. UK Journal of Pharmaceutical and Biosciences. 3(1): 18-22.
2. Afify AE, El-Beltagi HS, Aly AA, El-Ansary AE. Antioxidant enzyme activities and lipid peroxidation as biomarker for potato tuber stored by two essential oils from caraway and clove and its main component carvone and eugenol. Asian Pac J Trop Biomed 2012; 2(Suppl 2): S772-S780.
3. Aggarwal BB, Shishodia S. Molecular targets of dietary agents for prevention and therapy of cancer. Biochem Pharmacol 2006; 71(10): 1397-1421.
4. Ahmad I., M.A. Hanif, R. Nadeem, M.S. Jamil, M.S. Zafar. (2008). Nutritive evaluation of medicinal plants being used as condiments in South Asian Region. Journal of the Chemical Society of Pakistan. 30(3): 400-405.
5. Ali SM, Khan AA, Ahmed I, Musaddiq M, Ahmed KS, Polasa H, et al. Antimicrobial activities of eugenol and cinnamaldehyde against the human gastric pathogen *Helicobacter pylori*. Ann Clin Microbiol Antimicrob 2005; 4: 20.
6. Al-Maskri A.Y., M.A. Hanif, M.Y. Al-Maskari, A.S. Abraham, J.N. Al-sabahi, O. Al-Mantheri. (2011). Essential oil from *Ocimum basilicum* (Omani Basil): a desert crop. Natural product communications. 6(10): 1934578X1100601020.
7. Amla MH, Ertas M, Nitz S, Kollmannsberger H (2007). Chemical composition and content of essential oil from the bud of cultivated Turkish clove (*Syzygium aromaticum* L.). Bio-Res., 2(2): 265-269.
8. Arshad Z., M.A. Hanif, R.W.K. Qadri, M.M. Khan. (2014). Role of essential oils in plant diseases protection: a review. International Journal of Chemical and Biochemical Sciences. 6: 11-17.
9. Atsumi T, Fujisawa S, Tonosaki K. A comparative study of the antioxidant/prooxidant activities of eugenol and isoeugenol with various concentrations and oxidation conditions. Toxicol In Vitro 2005; 19(8): 1025-1033.
10. Ayoola GA, Lawore FM, Adelowotan T, Aibinu IE, Adenipekun E, et al., (2008) Chemical analysis and antimicrobial activity of the essential oil of *Syzygium aromaticum* (clove). African Journal of Microbiology Research 2(7): 162-166.
11. Banerjee S, Das, S. Anticarcinogenic effects of an aqueous infusion of cloves on skin carcinogenesis. Asian Pac J. Cancer Prev. 2005; 6(3): 304-308.
12. Banerjee S, Panda KC, Das, S. Clove (*Syzygium aromaticum* L.) A potential Chemopreventive agent for lung cancer. Carcinogenesis. 2006; 27(8):1645-1654.
13. Barbosa JD, Silva VB, Alves PB, Gumina G, Santos RL, Sousa DP, et al. Structure-activity relationships of eugenol derivatives against *Aedes aegypti* (Diptera: Culicidae) larvae. Pest Manag Sci 2012; 68(11): 1478-1483.
14. Bendre RS, Rajput JD, Bagul SD, Karandikar PS (2016) Outlooks on medicinal properties of eugenol and its synthetic derivatives. Natural Products Chemistry & Research 4(212): 2.
15. Bhuiyan MNI, Begum J, Nandi NC, Akter F (2010) Constituents of the essential oil from leaves and buds of clove (*Syzygium caryophyllum* (L.) Alston). African Journal of Plant Science 4(11): 451-454.
16. Campbell N. A. and J. B. Reece, Biology, Vol. 1, Erlangga, Jakarta, Indonesia, Eighth edition, 2012.
17. Chaiet al., Hajlaoui H, Zmantar T, Kahla-Nakbi, A.B., Rouabchia, M., Mahdouani, K. and Bakhrouf, A. The chemical composition and biological activity of essential oil, *Eugenia Caryophyllata* (*Syzygium aromaticum* L. Myrtaceae): a short review Phytotherapy Research. 2007; 21(6): 501-506.
18. Chami F, Chami N, Bennis S, Trouillas J, Remmal A. Evaluation of carvacrol and eugenol as prophylaxis and treatment of vaginal candidiasis in an immune suppressed rat model. J Antimicrob Chemother 2004; 54(5): 909-914.
19. Cock I.E., M. Cheesman. (2018). Plants of the genus *Syzygium* (Myrtaceae): A review on ethnobotany, medicinal properties and phytochemistry. Bioactive Compounds of Medicinal Plants. Ed Goyal MR, Ayeleso A. Apple Academic Press, USA.
20. Cortés-Rojas DF, de Souza CRF, Oliveira WP. Clove (*Syzygium aromaticum*): A precious spice. Asian Pac J Trop Biomed, 2014; 4(2): 90-6.
21. Daniel AN, Sartoretto SM, Schimidt G, Caparroz-Assef SM, Bersani-Amado CA, Cuman RK. Anti-inflammatory and antinociceptive activities of eugenol essential oil in experimental animal models. Rev Bras Farmacogn 2009; 19(1B): 212-217.
22. Danthu P, Penot E, Ranoarisoa KM, Rakotondravelo JC, Michel I, et al., (2014) The clove tree of Madagascar: a success story with an unpredictable future. Bois et forêts des tropiques 320 (2): 83-95.
23. Devi KP, Nisha SA, Sakthivel R, Pandian SK. Eugenol (an essential oil of clove) acts as an antibacterial agent against *Salmonella typhib* disrupting the cellular membrane. J Ethnopharmacol 2010; 130(1): 107-115.

24. Doleželová P, Mácová S, Plhalová L, Pištěková V, Svobodová Z. The acute toxicity of clove oil to fish *Danio rerio* and *Poecilia reticulata*. *Acta Vet Brno* 2011; 80(3): 305-308.

25. Dorman H, S.G. Deans. (2000). Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *Journal of applied microbiology*. 88(2): 308-316.

26. Dudonné S, Vitrac X, Coutière P, Woillez M, Mérillon JM. Comparative study of antioxidant properties and total phenolic content of 30 plant extracts of industrial interest using DPPH, ABTS, FRAP, SOD, and ORACassays. *J Agric Food Chem* 2009; 57(5): 1768-1774.

27. Eamsobhana P, Yoolek A, Kongkaew W, Lerdthusnee K, Khlaimanee N, Parsartvit A, et al. Laboratory evaluation of aromatic essential oils from thirteen plant species as candidate repellents against *Leptotrombiculidium chiggers* (Acar: Trombiculidae), the vector of scrub typhus. *Exp Appl Acarol* 2009; 47(3): 257-262.

28. Evert R. F., Esau's Plant Anatomy: Meristems, Cells, and Tissues of the Plant Body: Heir Structure, Function, and Development, John Wiley & Sons, Madison, WI, USA, Third edition, 2006.

29. Filho GA, Cesar JO, Ramos JV. [Cravo from India]. Itabuna: CEPLAC; 2013. [Online] Available from: <http://www.ceplac.gov.br/radar.htm>. [Accessed on 21st April, 2013]. Portuguese.

30. Fu Y, Zu Y, Chen L, Shi X, Wang Z, Sun S, et al. Antimicrobial activity of clove and rosemary essential oils alone and in combination. *Phytother Res* 2007; 21(10): 989-994.

31. Garg A, Singh S. Enhancement in antifungal activity of eugenol in immunosuppressed rats through lipid nanocarriers. *Colloids Surf B Biointerfaces* 2011; 87(2): 280-288.

32. Ghadermazi R, J. Keramat, S. Goli. (2017). Antioxidant activity of clove (*Eugenia caryophyllata* Thunb), oregano (*Origanum vulgare* L) and sage (*Salvia officinalis* L) essential oils in various model systems. *International Food Research Journal*. 24(4): 1628.

33. Ghelardini C, Galeotti N, Di Cesare Mannelli L, Mazzanti G, Bartolini A. Local anaesthetic activity of β -caryophyllene 11. *Farmaco*. 2001; 56: 387-389.

34. Ghosh R, Nadiminty N, Fitzpatrick JE, Alworth WL, Slaga TJ, Kumar AP. Eugenol causes melanoma growth suppression through inhibition of E2F1 transcriptional activity. *J Biol Chem* 2005; 280(7): 5812-5819.

35. Goni, P, López P, Sánchez C, Gómez Lus R, Becerril R, et al., (2009) Antimicrobial activity in the vapour phase of a combination of cinnamon and clove essential oils. *Food chemistry* 116(4): 982-989.

36. Guénette SA, Ross A, Marier JF, Beaudry F, Vachon P. Pharmacokinetics of eugenol and its effects on thermal hypersensitivity in rats. *Eur J Pharmacol* 2007; 562(1-2): 60-67.

37. Gulcin I, Sat IG, Beydemir S, Elmastas M, Kufrevioglu OI. Comparison of antioxidant activity of clove (*Eugenia caryophyllata* Thunb) buds and lavender (*Lavandula stoechas* L.). *Food Chemistry*. 2004; 87: 393-400.

38. Gürçin İ. Antioxidant activity of eugenol: a structure-activity relationship study. *J Med Food* 2011; 14(9): 975-985.

39. Hanif M.A., M.Y. Al-Maskari, A. Al-Maskari, A. Al-Shukaili, A.Y. Al-Maskari, J.N. Al-Sabahi. (2011). Essential oil composition, antimicrobial and antioxidant activities of unexplored Omani basil. *Journal of Medicinal Plants Research*. 5(5): 751-757.

40. Hanif M.A., S. Nisar, G.S. Khan, Z. Mushtaq, M. Zubair, Essential Oils. In *Essential Oil Research*, Springer: 2019; pp 3-17.

41. Healthcare T. PDR for herbal medicines. 4th ed. Montvale: Thomson Healthcare; 2004.

42. Hekimoğlu MA, Ergun M. Evaluation of clove oil as anaesthetic agent in fresh water angelfish, *Pterophyllum scalare*. *Pak J Zool* 2012; 44(5): 1297-1300.

43. Hill LE, Gomes C, Taylor TM. Characterization of beta-cyclodextrin inclusion complexes containing essential oils (trans-cinnamaldehyde, eugenol, cinnamon bark, and clove bud extracts) for antimicrobial delivery applications. *LWT-Food Sci Technol* 2013; 51(1): 86-93.

44. Huang Y, Ho SH, Lee HC, Yap YL (2002). Insecticidal properties of eugenol, isoeugenol and methyleugenol and their effects on nutrition of *Sitophilus zeamais* Motsch. (Coleoptera: Curculionidae) and *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *J. Stored Prod. Res.*, 38: 403-412.

45. Hussain N, Trak Th. Medicinal Virtues and Phytochemical Constituents of Some of World Journal of Pharmaceutical Research Medicinal Virtues and Phytochemical Constituents of, October, 2019.

46. International Organization for Standardization. Oil of clover leaf [Syzygium aromaticum (Linnaeus) Merril and Perry, syn. *Eugeniacyano* phyllus (Sprengel) Bullock and S.Harrison]. ISO-Directive 3141/1997, Geneva, Switzerland, 2002.

47. Jain A. International Journal of Scientific Research Therapeutic and Medicinal Uses Of Lavanga-A Review Ayurveda Dr. Nidhi Garg Th. 2019; 10: 8-10.

48. Javahery S, Nekobin H, Moradlu AH. Effect of anaesthesia with clove oil in fish (review). *Fish Physiol Biochem* 2012; 38(6): 1545-1552.

49. Javed S, A.A. Shahid, M.S. Haider, A. Umeera, R. Ahmad, S. Mushtaq. (2012). Nutritional, phytochemical potential and pharmacological evaluation of *Nigella Sativa* (Kalonji) and *Trachyspermum Ammi* (Ajwain). *Journal of Medicinal Plants Research*. 6(5): 768-775.

50. Jirovetz L, Buchbauer G, Stoilova I, Stoyanova A, Krastanov A, Schmidt E. Chemical composition and antioxidant properties of clover leaf essential oil. *J Agric Food Chem* 2006; 54(17): 6303-6307.

51. Kafle L, Shih CJ. Toxicity and repellency of compounds from clove (*Syzygium aromaticum*) to red imported fire ants *Solenopsis invicta* (Hymenoptera: Formicidae). *J Econ Entomol* 2013;106(1): 131-135.
52. Kamatou GP, Vermaak I, Viljoen AM. Eugenol--from the remote Maluku Islands to the international market place: a review of a remarkable and versatile molecule. *Molecules* 2012;17(6): 6953-6981.
53. Kamel C, Hafedh H, Tarek Z, Amel BKN, Mahmoud R, Kacem M, Amina B (2007). The chemical composition and biological activity of clove essential oil, *Eugenia caryophyllata* (*Syzygium aromaticum* L. Myrtaceae): A short review. *Phytother. Res.*, 21(6): 501-506.
54. Khalil AA, Rahman UU, Khan MR, Sahar A, Mehmood T, Khan M. Essential oil eugenol: Sources, extraction techniques and nutraceutical perspectives. *RSC Adv*, 2017; 7(52): 32669-81.
55. Khan M.M., M. Iqbal, M.A. Hanif, M.S. Mahmood, S.A. Naqvi, M. Shahid, M.J. Jaskani. (2012). Antioxidant and antipathogenic activities of citrus peel oils. *Journal of Essential Oil Bearing Plants*. 15(6): 972-979.
56. Körfers A, Sun Y. *Caryophylli flos- Gewürznelken-Ding X. Traditionelle Chinesische Medizin-Arzneidrogen und Therapie*, Wissenschaftliche Verlagsgesellschaft mbH Stuttgart Germany 2009, 397-398.
57. Kridati E. M., E. Prihastanti, and S. Haryanti, "Yield of essential oils and organ diameter and size of *Foeniculum vulgare* mill cells cultivated in semarang regency and Salatiga city," *Bulletin of Anatomy and Physiology*, vol. 20, no. 1, pp. 1-17, 2012.
58. Laitupa F. and H. Susane, Utilization of Eugenol from Clove Oil to Overcome Rancidity in Coconut Oil, Diponegoro University, Semarang, Indonesia, 2010.
59. Lee KG, Shibamoto T (2001). Antioxidant property of aroma extract isolated from clove buds [*Syzygium aromaticum* (L.) Merr. Et Perry]. *Food Chem.*, 74: 443-448.
60. Li HY, Lee BK, Kim JS, Jung SJ, Oh SB. Eugenol inhibits ATP-induced P2X currents in trigeminal ganglion neurons. *Korean J Physiol Pharmacol* 2008; 12(6): 315-321.
61. Lopez P, Sanchez C, Batle B, Nerin C. Solid and vapour phase antimicrobial activities of six essential oils: susceptibility of selected food borne bacterial and fungal strains. *Journal of Agriculture and Food Chemistry*. 2005; 53: 6338-6346.
62. Luthfi M., Management of Clove Bud Harvest (*Syzygium aromaticum* L.) at Kebun Branggah Banaran: Pt. Perkebunan Tjengkeh, Blitar, East Java, Bogor Agriculture Institute, Bogor, Indonesia, 2017.
63. Milind P., K. Deepa. (2011). Clove: a champion spice. *Int J Res Ayurveda Pharm.* 2(1): 47-54.
64. Mittal M., N. Gupta, P. Parashar, V. Mehra, M. Khatri. (2014). Phytochemical evaluation and pharmacological activity of *Syzygium aromaticum*: a comprehensive review. *Int J Pharm Pharm Sci.* 6(8): 67-72.
65. Muruganadan S, Srinivasan K, Chandra S, Tandan SK, Lal J, Raviprakash V (2001). Anti-inflammatory activity of *Syzygium cuminibark*. *Fitoterapia*, 72: 369-375.
66. Nam H, Kim MM. Eugenol with antioxidant activity inhibits MMP-9 related to metastasis in human fibrosarcoma cells. *Food Chem Toxicol* 2013;55: 106-112.
67. Nassar MI (2006). Flavonoid triglycosides from the seeds of *Syzygium aromaticum*. *Carbohydrate Res.*, 341: 160-163.
68. Nuraniza, B. P. Lapanporo, and Y. Arman, "Edible oil quality test based on changes in light polarization angles using semiautomatic polarimeter," *ISSN Journal*, vol. 1, no. 2, pp. 87-91, 2013.
69. Oliveira RA, Oliveira FF, Sacramento CK. [Essential oils: prospects for agribusiness spices in Bahia]. *Bahia Agric* 2007; 8(1): 46-48. Portuguese.
70. Oliveira RA, Reis TV, Sacramento CK, Duarte LP, Oliveira FF. Volatile chemical constituents of rich spices in eugenol. *Rev Bras Farmacognosia* 2009; 19(3): 771-775.
71. Overly KR. Microwave-Assisted Isolation of Eugenol from Cloves. *J Chem Educ.*, 2019; 96(11): 2665-7.
72. Pambudi N. K., S. Sunarta, and Kasmudjo , Effect of Variation in Plant Age and Leaves on Leaves and Quality of eucalyptus (*Eucalyptus Alba Reinw. Ex. BI*): Essay, Gadjah Mada University, Yogyakarta, Indonesia, 2016.
73. Park IK, Shin SC. Fumigant activity of plant essential oils and components from garlic (*Allium sativum*) and clove bud (*Eugenia caryophyllata*) oils against the Japanese termite (*Reticulitermes speratus Kolbe*). *J Agric Food Chem* 2005; 53(11): 4388-4392.
74. Park M.-J., K.-S. Gwak, I. Yang, W.-S. Choi, H.-J. Jo, J.-W. Chang, E.-B. Jeung, I.-G. Choi. (2007). Antifungal activities of the essential oils in *Syzygium aromaticum* (L.) Merr. Et Perry and *Leptospermum petersonii* Bailey and their constituents against various dermatophytes. *The Journal of Microbiology*. 45(5): 460-465.
75. Pérez-Conesa D, McLandsborough L, Weiss J. Inhibition and inactivation of *Listeria monocytogenes* and *Escherichia coli* O157:H7 colony biofilms by micellar-encapsulated eugenol and carvacrol. *J Food Prot* 2006;69 (12): 2947-2954. Diego Francisco Cortés-Rojas et al./Asian Pac J Trop Biomed 2014; 4(2): 90-96 96
76. Pérez-Jiménez J, Neveu V, Vos F, Scalbert A. Identification of the 100richest dietary sources of polyphenols: an application of the phenol-explorer database. *Eur J Clin Nutr* 2010; 64(Suppl 3): S112-S120.
77. Pino JA, Marbot R, Agüero J, Fuentes V (2001). Essential oil from buds and leaves of clove (*Syzygium aromaticum* (L.) Merr. et Perry) grown in Cuba. *Journal of Essential Oil Research* 13(4): 278-279.
78. Pinto E, Valer-Silva L, Cavaleiro C, Salgueiro E. Antifungal activity of clove essential oil from *Syzygium aromaticum* on *candida aspergillus* and dermatophyte species. 2009; 58:1454-1462.

79. Pleasant, Barbara (2011) "All About Growing Strawberries". Mother Earth News (248): 23-25.
80. Prasad RC, Herzog B, Boone B, Sims L, Waltner-Law L. An extract of *Syzygium aromaticum* represses genes encoding hepatic gluconeogenic enzymes. *J Ethnopharmacol.* 2005; 96(1-2):295-301.
81. Pujiarti R., Y. Ohtani, and H. Ichiura, "Physicochemical properties and chemical compositions of *Melaleuca leucadendron* leaf oils taken from the plantations in Java, Indonesia," *Journal of Wood Science*, vol. 57, no. 5, pp. 446-451, 2011.
82. Raina VK, Srivastava SK, Aggarwal KK, Syamasundar KV, Sushil K(2001). Essential oil composition of *Syzygium aromaticum* leaf from Little Andaman, India. *Flav. Fra. J.*, 16(5): 334-336.
83. Rana IS, Rana AS, Rajak RC. Evaluation of antifungal activity in essential oil of the *Syzygium aromaticum* (L.) by extraction, purification and analysis of its main component eugenol. *Braz J Microbiol* 2011; 42(4): 1269-1277.
84. Razafimamonjison G., M. Jahiel, P. Ramanoelina, F. Fawbush, and P. Danthu, "Effect of phenological stages on yield and composition of essential oils of *Syzygium aromaticum* buds from Madagascar," *International Journal of Basic and Applied Sciences*, vol. 2, no. 4, pp. 312-318, 2013.
85. Rukmana R. and H. Yudirachman, *Luckily Exorbitant from Clove Agribusiness*, Lily Publisher, Yogyakarta, Indonesia, 2016.
86. Runtunuwu S. D., R. Mamarimbings, P. Tumewu, and R. M. N. Rengkung, "Effect of paclobutrazol on the quality of clove (*Syzygium aromaticum*L)," *Journal of Bioslogos*, vol. 6, no. 2, pp. 33-41, 2016.
87. Sallato BV, Torres R, Zoffoli JP, Latorre BA (2007) Effect of boscalid on postharvest decay of strawberry caused by *Botrytis cinerea* and *Rhizopus stolonifer*. *Spanish journal of agricultural research* 5(1): 67-78.
88. Schönfelder I, Schönfelder P. *Syzygiumaromaticum* (L.) MERR & PERRY (Eugeniacyophyllata THUNB Jambosacaryophyllus (SPRENG.) (NIEDENZU). Das neueHandbuch der Heilpflanzen, Kosmos-Verlag, Stuttgart, Germany, 2004, 431-432.
89. Semangun H. and F. F. Karwur, *Cloves: History, Cultivation, and Industry*, Gramedia, Jakarta, Indonesia, 2014.
90. Shafi PM, Rosamma MK, Jamil K, Reddy PS (2002). Antibacterial activity of *S. cumini* and *S. travancoricum* leaf essential oils. *Fitoterapia*, 73: 414-416.
91. Shan B, Cai YZ, Sun M, Corke H. Antioxidant capacity of 26spice extracts and characterization of their phenolic constituents. *J Agric Food Chem* 2005; 53(20): 7749-7759.
92. Shobana S., K.A. Naidu. (2000). Antioxidant activity of selected Indian spices. *Prostaglandins, Leukotrienes and Essential Fatty Acids (PLEFA)*. 62(2): 107-110.
93. Singh AK, Dhamanigi SS, Asad M. Anti-stress activity of hydro-alcoholic extract of *Eugenia caryophyllus* buds (clove). *Indian J. Pharmacol.* 2009; 41:28-31
94. Slamenová D, Horváthová E, Wsólová L, Sramková M, Navarová J. Investigation of anti-oxidative, cytotoxic, DNA-damaging and DNA-protective effects of plant volatiles eugenol and borneol in human-derived HepG2, Caco-2and VH10cell lines. *Mutat Res* 2009; 677(1-2): 46-52.
95. Sofia P.K., R. Prasad, V.K. Vijay, A.K. Srivastava. (2007). Evaluation of antibacterial activity of Indian spices against common foodborne pathogens. *International journal of food science & technology*. 42(8): 910-915.
96. Sritabutra D, Soonwera M, Waltanachanobon S, Poungjai S. Evaluation of herbal essential oil as repellents against *Aedes aegypti* (L.)and *Anopheles dirus* Peyton & Harrion. *Asian Pac J Trop Biomed* 2011; 1(Suppl 1): S124-S128.
97. Srivastava AK, Srivastava SK, Syamsundar KV (2003). Bud and leaf essential oil composition of *Syzygium aromaticum* from India and Madagascar. *Flav. Fra. J.*, 20(1): 51-53.
98. Tajuddin, Ahmed S, Latif A, Qasmi IA. Effect of 50% ethanolic extract of *Syzygium aromaticum* (L.) Merr. & Perry. Clove on sexual behaviour of normal male rats. *BMC Complement Altern.Med.* 2004; 4: 17.
99. Thangaselvabai T, Kennedy R, Joshua J, Jayasekar M. Clove (*Syzygium Aromaticum*) -The Spicy Flower Bud of Significance - A Review. *Agric Rev.*, 2010; 31(1): 40-47.

Submit your next manuscript to ScienceFrontier and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Research which is freely available for redistribution
- Submit your manuscript at: <https://sciencefrontier.org/submit-manuscript?e=2>



© The Author(s) 2026. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license,